

Array and Queries

Given an array, you are asked to perform a number of queries and divide the array into what are called, *beautiful* subsequences.

The array A has length n . A function $f(A)$ is defined to be a minimal possible x , such that it's possible to divide array A into x *beautiful* subsequences. Note that each element of an array should belong to exactly one subsequence, and subsequence does not necessarily need to be consecutive.

A subsequence S with length len is called *beautiful* if and only if:

- $len = 1$ or
- Let S' be a sorted version of S . It must hold that $S'_i = S'_{i+1} - 1$ for every $i \in [1, len - 1]$.

For instance, if $A = [1, 2, 3, 4, 3, 5]$, $f(A)$ would be 2. Because, you can divide A into 2 *beautiful* subsequences either like $[1, 2, 3]$ and $[4, 3, 5]$ or like $[1, 2, 3, 4, 5]$ and $[3]$.

You have to answer q queries. Each query is of the type:

- ***id val***: you need to change a value of A_{id} to val , i.e. $A_{id} = val$. Here *id* is 1 – *indexed*.

After each query, for the value of $f(A)$, let's denote that value as ans_i , where i indicates the i^{th} query.

You need to find $\sum_{i=1}^q i \times ans_i$ modulo $(10^9 + 7)$.

Input Format

The first line contains a single integer n , representing the length of array A .

The next line contains the array A given as space-separated integers.

The next line contains a single integer q , representing the number of queries.

Each of the q lines contain two integers *id* and *val*, which is described above.

Constraints

- $1 \leq n, q \leq 3 \times 10^5$
- $1 \leq A_i \leq 10^9$
- $1 \leq id \leq n$
- $1 \leq val \leq 10^9$

Output Format

Print the required answer in one line.

Sample Input 0

```

5
2 2 1 1 1
2
3 2
5 5

```


Sample Output 0

11

Explanation 0

The initial array A is $[2, 2, 1, 1, 1]$

- After 1^{st} query the array becomes $[2, 2, 2, 1, 1]$ this can be divided into **3** subsequences as $[2, 1]$, $[2, 1]$ and $[2]$.
- After 2^{nd} query the array becomes $[2, 2, 2, 1, 5]$ this can be divided into **4** subsequences as $[2, 1]$, $[2]$, $[2]$ and $[5]$.



Hence, calculating $\sum i \times ans_i$ we get

$$1 \times 3 + 2 \times 4 \Rightarrow 11$$

Sample Input 1

```

2
3 3
3
2 4
1 5
2 2

```

Sample Output 1

9

Explanation 1

The initial array A is $[3, 3]$

- After 1^{st} query the array becomes $[3, 4]$ this can be divided into **1** subsequence as $[3, 4]$.
- After 2^{nd} query the array becomes $[5, 4]$ this can be divided into **1** subsequence as $[5, 4]$.
- After 3^{rd} query the array becomes $[5, 2]$ this can be divided into **2** subsequences as $[5]$ and $[2]$.

Hence, calculating $\sum i \times ans_i$ we get

$$1 \times 1 + 2 \times 1 + 3 \times 2 \Rightarrow 9$$