

John and GCD list

John is new to Mathematics and does not know how to calculate [GCD](#) of numbers. So he wants you to help him in a few GCD calculations. John has a list A of numbers, indexed 1 to N . He wants to create another list B having $N+1$ numbers, indexed from 1 to $N+1$, and having the following property:

$$\text{GCD}(B[i], B[i+1]) = A[i], \forall 1 \leq i \leq N$$

As there can be many such lists, John wants to know the list B in which sum of all elements is minimum. It is guaranteed that such a list will always exist.

Input Format

The first line contains an integer T , i.e., the number of the test cases. T testcases follow.

The first line of each test case contains an integer N , i.e., the number of elements in the array.

The second line of each test case contains N space separated integers that denote the elements of the list A .

Output Format

For each test case, print in a new line the list B such that each element is separated by a single space.

Constraints

$$1 \leq T \leq 10$$

$$2 \leq N \leq 10^3$$

$$1 \leq A[i] \leq 10^4$$

$$1 \leq B[i]$$

Sample Input

```
2
3
1 2 3
3
5 10 5
```

Sample Output

```
1 2 6 3
5 10 10 5
```

Explanation

For the first testcase,

```
GCD(1,2) = 1
GCD(2,6) = 2
GCD(6,3) = 3
sum = 1+2+6+3 = 12 which is minimum among all possible list B
```

For the second testcase,

$$\text{GCD}(5, 10) = 5$$

$$\text{GCD}(10, 10) = 10$$

$$\text{GCD}(10, 5) = 5$$

$$\text{sum} = 5 + 10 + 10 + 5 = 30 \text{ which is the minimum among all possible list B}$$