

Alice thinks of a non-decreasing sequence of non-negative integers and wants Bob to guess it by providing him the set of all its **K**-sums with repetitions.

What is this? Let the sequence be $\{A[1], A[2], \dots, A[N]\}$ and **K** be some positive integer that both Alice and Bob know. Alice gives Bob the set of all possible values that can be generated by this - $A[i_1] + A[i_2] + \dots + A[i_K]$, where $1 \leq i_1 \leq i_2 \leq \dots \leq i_K \leq N$. She can provide the values generated in any order she wishes to. Bob's task is to restore the initial sequence.

Consider an example. Let **N = 3** and **K = 2**. The sequence is $\{A[1], A[2], A[3]\}$. The sequence of its **2**-sums with repetitions is $\{A[1] + A[1], A[1] + A[2], A[1] + A[3], A[2] + A[2], A[2] + A[3], A[3] + A[3]\}$. But its elements could be provided in any order. For example any permutation of **{2, 3, 4, 4, 5, 6}** corresponds to the sequence **{1, 2, 3}**.

Input Format

The first line of the input contains an integer **T** denoting the number of test cases.
The description of **T** test cases follows.
The first line of each test case contains two space separated integers **N** and **K**.
The second line contains the sequence **S_i** of all **K**-sums with repetitions of the sequence Alice initially thought of.

Constraints

- $1 \leq T \leq 10^5$
- $1 \leq N \leq 10^5$
- $1 \leq K \leq 10^9$
- $2 \leq S_i \leq 10^{18}$

Note

The total number of elements in any input sequence does not exceed **10⁵**
Each element of each input sequence is non-negative integer not exceeding **10¹⁸**.
Each input sequence is a correct sequence of all **K**-sums with repetitions of some non-decreasing sequence of non-negative integers.

Output Format

For each test case, output a single line containing the space separated list of elements of the non-decreasing sequence Alice thinks of. If there are several possible outputs you can output any of them.

Sample Input 0

```
3
1 3
3
2 2
```

```
12 34 56
3 2
2 3 4 4 5 6
```

Sample Output 0

```
1
6 28
1 2 3
```

Explanation 0

Sample case #00: When $N = 1$ and $K = 3$ the only K -sum is $S[1] = 3 * A[1]$. Hence $A[1] = S[1] / 3 = 3 / 3 = 1$.

Sample case #01: Since $6 + 6 = 12$, $6 + 28 = 34$, $28 + 28 = 56$, then Alice indeed could think of the sequence $\{6, 28\}$.

Sample case #02: It corresponds to the example in the problem statement.