Xoring Ninja

HackerRank

An XOR operation on a list is defined here as the *xor* (\oplus) of all its elements (e.g.: $XOR(\{A, B, C\}) = A \oplus B \oplus C$).

The XorSum of set arr is defined here as the sum of the XORs of all non-empty subsets of arr known as arr'. The set arr' can be expressed as:

$$XorSum(arr) = \sum_{i=1}^{2^{n}-1} XOR(arr'_{i}) = XOR(arr'_{1}) + XOR(arr'_{2}) + \dots + XOR(arr'_{2^{n}-2}) + XOR(arr'_{2^{n}-1})$$

For example: Given set $arr = \{n_1, n_2, n_3\}$

- The set of possible non-empty subsets is: $arr' = \{\{n_1\}, \{n_2\}, \{n_3\}, \{n_1, n_2\}, \{n_1, n_3\}, \{n_2, n_3\}, \{n_1, n_2, n_3\}\}$
- The XorSum of these non-empty subsets is then calculated as follows: $XorSum(arr) = n_1 + n_2 + n_3 + (n_1 \oplus n_2) + (n_1 \oplus n_3) + (n_2 \oplus n_3) + (n_1 \oplus n_2 \oplus n_3)$

Given a list of n space-separated integers, determine and print $XorSum \% (10^9 + 7)$.

For example, $arr = \{3,4\}$. There are three possible subsets, $arr' = \{\{3\}, \{4\}, \{3,4\}\}$. The XOR of arr'[1] = 3, of arr'[2] = 4 and of $arr[3] = 3 \oplus 4 = 7$. The XorSum is the sum of these: 3 + 4 + 7 = 14 and $14 \% (10^9 + 7) = 14$.

Note: The cardinality of powerset(n) is 2^n , so the set of non-empty subsets of set arr of size n contains $2^n - 1$ subsets.

Function Description

Complete the *xoringNinja* function in the editor below. It should return an integer that represents the XorSum of the input array, modulo $(10^9 + 7)$.

xoringNinja has the following parameter(s):

• arr: an integer array

Input Format

The first line contains an integer T, the number of test cases.

Each test case consists of two lines:

- The first line contains an integer n, the size of the set arr.
- The second line contains n space-separated integers arr[i].

Constraints

 $egin{arred} &1 \leq T \leq 5 \ &1 \leq n \leq 10^5 \ &0 \leq arr[i] \leq 10^9, \ &1 \leq i \leq n \end{aligned}$

Output Format

For each test case, print its $XorSum \% (10^9 + 7)$ on a new line. The i^{th} line should contain the output for the i^{th} test case.

Sample Input 0

Sample Output 0

12

Explanation 0

The input set, $S = \{1, 2, 3\}$, has 7 possible non-empty subsets: $S' = \{\{1\}, \{2\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 3\}, \{1, 2, 3\}\}.$

We then determine the XOR of each subset in S^{\prime} :

 $XOR(\{1\}) = 1$ $XOR(\{2\}) = 2$ $XOR(\{3\}) = 3$ $XOR(\{1,2\}) = 1 \oplus 2 = 3$ $XOR(\{2,3\}) = 2 \oplus 3 = 1$ $XOR(\{1,3\} = 1 \oplus 3 = 2$ $XOR(\{1,2,3\} = 1 \oplus 2 \oplus 3 = 0$

Then sum the results of the XOR of each individual subset in S', resulting in XorSum = 12 and $12~\%~(10^9 + 7) = 12$.

Sample Input 1

2 4 1 2 4 8 5 1 2 3 5 100

Sample Output 1

120 1648